

The correspondence between a plane curve and its complement

By *Jérémy Blanc* at Saint-Martin d'Hères

Abstract. Given two irreducible curves of the plane which have isomorphic complements, it is natural to ask whether there exists an automorphism of the plane that sends one curve on the other.

This question has a positive answer for a large family of curves and H. Yoshihara conjectured that it is true in general. We exhibit counterexamples to this conjecture, over any ground field. In some of the cases, the curves are isomorphic and in others not; this provides counterexamples of two different kinds.

Finally, we use our construction to find the existence of surprising non-linear automorphisms of affine surfaces.

1. Introduction

In this article, \mathbb{K} is any field, and all surfaces are algebraic affine or projective surfaces, defined over \mathbb{K} .

1.1. The conjecture. To any irreducible curve $C \subset \mathbb{P}^2 = \mathbb{P}_{\mathbb{K}}^2$ we can associate its complement, the affine surface $\mathbb{P}^2 \setminus C$ (such affine surfaces have been a subject of research for many years, see [2], [3], [9], [6], [7], [4], [5], ...). If two such curves C, D are projectively equivalent—i.e. if some automorphism of the projective plane \mathbb{P}^2 sends C on D —then clearly $\mathbb{P}^2 \setminus C$ is isomorphic to $\mathbb{P}^2 \setminus D$. It is natural to ask whether the converse is true. In 1984, Hisao Yoshihara made the following conjecture.

Conjecture 1.1 ([11]). *Let $C \subset \mathbb{P}_{\mathbb{K}}^2$ be an irreducible curve and assume that \mathbb{K} is algebraically closed of characteristic 0. Suppose that $\mathbb{P}^2 \setminus C$ is isomorphic to $\mathbb{P}^2 \setminus D$ for some curve D . Then C and D are projectively equivalent.*

In [11], it was proved that the conjecture is true for a large family of curves C . We briefly recall these results in Section 2, and extend some of them to any field \mathbb{K} . Then, we

The author acknowledges support from the Swiss national science foundation.

provide a family of counterexamples to the conjecture, over any field \mathbb{K} , and prove the following result.

Theorem 1. *For any field \mathbb{K} with more than two elements, there exist two curves $C, D \subset \mathbb{P}^2_{\mathbb{K}}$, irreducible over the algebraic closure of \mathbb{K} , such that the following two assertions are true:*

- (1) *The affine surfaces $\mathbb{P}^2 \setminus C$ and $\mathbb{P}^2 \setminus D$ are isomorphic.*
- (2) *No automorphism of \mathbb{P}^2 sends C on D .*

Furthermore, there are examples where C and D are isomorphic and examples where they are not.

Observe that Theorem 1 yields the existence of isomorphic affine surfaces having a projective completion in isomorphic projective surfaces by irreducible non-isomorphic curves. Such examples were, as far as we are aware, not known before.

Recall that a curve C is of type I if there exists some point $a \in C$ such that $C \setminus a$ is isomorphic to the affine line. The problem stated above is related to another conjecture, namely:

Conjecture 1.2 ([12], page 101). *If $C \subset \mathbb{P}^2_{\mathbb{C}}$ is an irreducible curve, which is neither of type I nor a nodal cubic curve, then any automorphism of $\mathbb{P}^2 \setminus C$ extends to an automorphism of \mathbb{P}^2 .*

The construction we provide to prove Theorem 1 will also provide counterexamples to Conjecture 1.2, extending furthermore the possibilities for the base field.

Theorem 2. *Assume that the characteristic of \mathbb{K} is not 2. Then, there exists a curve $C \subset \mathbb{P}^2_{\mathbb{K}}$, irreducible over the algebraic closure of \mathbb{K} , of degree 39, that is not of type I, and there exists an automorphism of $\mathbb{P}^2_{\mathbb{K}} \setminus C$ that does not extend to $\mathbb{P}^2_{\mathbb{K}}$.*

1.2. The construction. Here, we briefly describe our construction, which will be explained more precisely in Section 3. We denote by $\Delta \subset \mathbb{P}^2$ the union of three general lines and choose two quartics Γ_1, Γ_2 that intersect Δ in a particular manner. We construct a birational morphism $\pi : X \rightarrow \mathbb{P}^2$ that is a sequence of blow-ups of points that belong, as proper or infinitely near points, to $\Delta \cap \Gamma_1$ or $\Delta \cap \Gamma_2$. Then, we find a reducible curve $R \subset \pi^{-1}(\Delta)$ such that for $i = 1, 2$, the curve $R \cup \widetilde{\Gamma}_i$ is contractible via a birational morphism $\eta_i : X \rightarrow \mathbb{P}^2$ (where $\widetilde{\Gamma}_i$ is the strict transform of Γ_i on X). The birational map $\varphi = \eta_1 \circ \eta_2^{-1}$ restricts to an isomorphism from $\mathbb{P}^2 \setminus \eta_2(\widetilde{\Gamma}_1)$ to $\mathbb{P}^2 \setminus \eta_1(\widetilde{\Gamma}_2)$.

$$(1) \quad \begin{array}{ccc} & & \mathbb{P}^2 \\ & \nearrow^{\eta_1} & \uparrow \varphi \\ \mathbb{P}^2 & \xleftarrow{\pi} X & \\ & \searrow_{\eta_2} & \downarrow \varphi \\ & & \mathbb{P}^2 \end{array}$$

In our construction, the curves Γ_1 and Γ_2 depend on parameters. For general values of these parameters, the curves $\eta_2(\widetilde{\Gamma}_1)$ and $\eta_1(\widetilde{\Gamma}_2)$ are not projectively equivalent, which yields the proof of Theorem 1. For special values of the parameters, there exists some auto-

morphism ψ of \mathbb{P}^2 that sends $\eta_2(\widetilde{\Gamma}_1)$ on $\eta_1(\widetilde{\Gamma}_2)$. Thus, $\varphi \circ \psi^{-1}$ is an automorphism of $\mathbb{P}^2 \setminus \eta_1(\widetilde{\Gamma}_2)$ that does not extend to an automorphism of \mathbb{P}^2 , which proves Theorem 2.

1.3. Outline of this article. In Section 2, we prove that Conjectures 1.1 and 1.2 are true for “most” kinds of curves. In Section 3, we describe precisely the construction announced in (1.2). Finally in Section 4 we prove that neither of the curves constructed is of type I, and decide when the curves obtained are projectively equivalent or isomorphic, which yields the proofs of Theorems 1 and 2.

1.4. Acknowledgements. The author presented the results of this article in Dijon and Genève. He would like to express his sincere gratitude to the members of these institutes for valuable questions which helped him to improve the exposition of this paper, with special thanks to Adrien Dubouloz and Thierry Vust.

2. Cases in which the conjectures are true

In this section, we prove that the conjectures are true for most curves, and recall some classical results. We will denote the algebraic closure of \mathbb{K} by $\overline{\mathbb{K}}$.

Definition 2.1. We say that a birational morphism $\chi : S \rightarrow \mathbb{P}^2$ is an *n-tower resolution* of a curve $C \subset \mathbb{P}^2$ if

- (1) the map χ decomposes as $\chi = \chi_m \circ \chi_{m-1} \circ \cdots \circ \chi_1$, for some integer $m \geq 0$, where χ_i is the blow-up of a point p_i and $\chi_{i-1}(p_i) = p_{i-1}$ for $i = 2, \dots, m$;
- (2) the strict transform of the curve C on S is a curve that is smooth, irreducible over $\overline{\mathbb{K}}$, isomorphic to \mathbb{P}^1 , and of self-intersection n .

Note that if a curve admits an n -tower resolution, it admits an m -tower resolution for any $m \leq n$. Next, we remind the reader of a simple but useful lemma, obvious for the specialist.

Lemma 2.2. *Let $C \subset \mathbb{P}^2$ be a curve irreducible over $\overline{\mathbb{K}}$, and let $\psi : \mathbb{P}^2 \setminus C \rightarrow \mathbb{P}^2 \setminus D$ be an isomorphism, where D is some curve of \mathbb{P}^2 .*

Then, either ψ extends to an automorphism of \mathbb{P}^2 (and in particular C and D are projectively equivalent), or there exist two birational morphisms $\chi, \varepsilon : S \rightarrow \mathbb{P}^2$ satisfying the following conditions:

- (1) χ (respectively ε) is a (-1) -tower resolution of C (respectively of D).
- (2) χ is a minimal resolution of the indeterminacies of ψ and $\psi\chi = \varepsilon$.

Proof. In this proof, we consider our algebraic varieties over the field $\overline{\mathbb{K}}$, remembering that these are defined over the subfield \mathbb{K} . We extend ψ to a birational transformation $\overline{\psi}$ of $\mathbb{P}^2_{\overline{\mathbb{K}}}$, which is defined over the field \mathbb{K} . Then, there exists a birational morphism $\overline{\chi} : S_{\overline{\mathbb{K}}} \rightarrow \mathbb{P}^2_{\overline{\mathbb{K}}}$, also defined over \mathbb{K} , that is a minimal resolution of the indeterminacies of $\overline{\psi}$. We denote the birational morphism $\overline{\psi} \circ \overline{\chi}$ by $\overline{\varepsilon}$ and denote by E (respectively F) the set of irreducible curves of $S_{\overline{\mathbb{K}}}$ that are collapsed by $\overline{\chi}$ (respectively by $\overline{\varepsilon}$). Since ψ is an

isomorphism of $\mathbb{P}^2 \setminus C$ to $\mathbb{P}^2 \setminus D$, and under the assumption that $\bar{\psi}$ is not an automorphism of $\mathbb{P}_{\bar{\mathbb{K}}}^2$, the map $\bar{\psi}$ collapses exactly one irreducible curve of $\mathbb{P}_{\bar{\mathbb{K}}}^2$, which is the extension of C as $\bar{C} \subset \mathbb{P}_{\bar{\mathbb{K}}}^2$. This means that the set $F \setminus E$ consists of a single element, which is the strict transform of \bar{C} ; since the sets E and F have the same number of curves, the set $E \setminus F$ also consists of a single element. This element has to be the strict transforms on $S_{\bar{\mathbb{K}}}$ of the extension \bar{D} of the curve D .

The resolution of $\bar{\psi}$ by $\bar{\chi}$ and $\bar{\varepsilon}$ being minimal, every irreducible curve of $E \cap F$ has self-intersection ≤ -2 ; this implies that the strict transforms of \bar{C} and \bar{D} on $S_{\bar{\mathbb{K}}}$ are (-1) -curves, i.e. both are smooth, irreducible, isomorphic to \mathbb{P}^1 and of self-intersection -1 .

The fact that only one irreducible curve collapsed by $\bar{\chi}$ (respectively by $\bar{\varepsilon}$) has self-intersection -1 implies that $\bar{\chi}$ is a tower resolution of $\bar{C} \subset \mathbb{P}_{\bar{\mathbb{K}}}^2$ (respectively of $\bar{D} \subset \mathbb{P}_{\bar{\mathbb{K}}}^2$). Since the set of points blown-up by both morphisms is invariant under the action of $\text{Gal}(\bar{\mathbb{K}}/\mathbb{K})$, and since no two points belong to the same surface, each point is defined over \mathbb{K} . Consequently, reducing the ground field to \mathbb{K} , we find birational morphisms χ and ε that are (-1) -tower resolutions of C and D respectively. \square

Corollary 2.3. *Conjectures 1.1 and 1.2 are true for any base field \mathbb{K} and any curve $C \subset \mathbb{P}^2$, irreducible over $\bar{\mathbb{K}}$, that does not admit a (-1) -tower resolution.*

In particular, both conjectures are true if C is not rational or if C has more than two singular points over $\bar{\mathbb{K}}$. \square

The conjectures are thus true for a large family of curves. Among curves admitting a tower resolution, curves of type I or II are the most natural to deal with. We remind the reader of some results on this subject.

Definition 2.4. A curve $C \subset \mathbb{P}^2$ is of type I (respectively of type II) if there exists a point $a \in C$ (respectively a line $L \subset \mathbb{P}^2$) such that $C \setminus a$ (respectively $C \setminus L$) is isomorphic to the affine line.

Any curve of type II is of type I and it is difficult (but possible) to find curves of type I that are not of type II [10]. A curve is of type II if and only if it is the image of a line by an automorphism of $\mathbb{P}^2 \setminus L$, where L is a line [1]. Furthermore, any curve of type II admits an n -tower resolution, for some positive integer $n \geq 3$ [13]. The following result gives another evidence to Conjecture 1.1:

Proposition 2.5 ([11]). *Conjecture 1.1 is true, over any algebraically closed field of characteristic 0, if C is of type II.*

Finally, we recall that Conjecture 1.1 was proved in [11], Proposition 2.7, in the case of a nodal cubic curve, and that the group $\text{Aut}(\mathbb{P}^2 \setminus C)$ for this curve was studied by Wakabayashi and Yoshihara, see [12] and [8].

3. The construction

In this section, we describe precisely the construction announced in the introduction. First we describe the triangle Δ , its irreducible components and singular points. Take three

general lines of \mathbb{P}^2 , that form a triangle Δ , and choose the coordinates such that Δ has equation $xyz = 0$. We denote by $a = (1 : 0 : 0)$, $b = (0 : 1 : 0)$, $c = (0 : 0 : 1) \in \mathbb{P}^2$ the singular points of Δ and by L_{ab} (respectively L_{ac} , L_{bc}) the line through a and b . In particular, $\Delta = L_{ab} \cup L_{ac} \cup L_{bc}$.

Then, we briefly describe the two curves Γ_1 and Γ_2 , in simple words. In subsection 3.1, we will describe these curves using the points infinitely near to a and b . For any $\theta \in \mathbb{K}^*$, we write $p(\theta) = (\theta : 0 : 1) \in L_{ac} \setminus \{a, c\}$ and denote by Ω_θ the set of irreducible quartic curves of \mathbb{P}^2 that have multiplicity 3 at $p(\theta)$, that pass through a being tangent to L_{ab} and intersect L_{bc} only at the point b .

Let $\alpha, \beta \in \mathbb{K}^*$, $\alpha \neq \beta$, then Γ_1 is one curve of Ω_α and Γ_2 is the curve of Ω_β whose intersection with Γ_1 at the point b is as large as possible.

3.1. The points in the neighbourhoods of a and b . We now describe the intersection between the curves Γ_1 , Γ_2 and Δ , and construct the birational morphism $\pi : X \rightarrow \mathbb{P}^2$ announced in Section 1.2.

We construct π by a sequence of blow-ups of points that lie on the curves Γ_1 , Γ_2 , Δ . Taking some point x in a surface S , the blow-up $p_x : S' \rightarrow S$ gives a smooth surface S' . We denote by $E_x \subset S'$ the exceptional curve of x , which is equal to $(p_x)^{-1}(x)$. Then, p_x is an isomorphism of $S' \setminus E_x$ to $S \setminus x$. It is therefore natural, for any point $y \in S \setminus x$ and any curve $C \subset S \setminus x$, to denote once again the point $p_x^{-1}(y)$ by y and the curve $p_x^{-1}(C)$ by C . For any curve $C \subset S$ passing through x , the strict transform of C on S' will be denoted by \tilde{C} . After two (or more) blow-ups, we write $\tilde{C} = \tilde{\tilde{C}}$ to simplify the notation.

Our aim is to obtain the configuration of curves of Figure 2 on X . For this, we will blow-up the points $p(\alpha)$, $p(\beta)$, and points in the neighbourhoods of a and b .

Denote by a_1 the point in the first neighbourhood of a that belongs to the (strict transform of the) line L_{ab} , and by b_1 the point in the first neighbourhood of b that belongs to the line L_{bc} . For $i = 2, 3$, we call b_i the point in the first neighbourhood of b_{i-1} (and thus in the i -th neighbourhood of b) that belongs to the line L_{bc} . We denote by $\pi' : X' \rightarrow \mathbb{P}^2$ the blow-up of the points a , a_1 , b , b_1 , b_2 , b_3 , $p(\alpha)$ and $p(\beta)$. The configuration of the curves on X' and the decomposition of π' are described in Figure 1.




Figure 1. The configuration of the special curves on the surface X' . Two curves are connected by an edge if their intersection is positive (and here equal to 1). The positive intersections with $\tilde{\Gamma}_1$ and $\tilde{\Gamma}_2$ are in square brackets.

On the surface X' , (the strict pull-back of) any curve of Ω_α has self-intersection 1, and its intersection with $E_{p(\alpha)}$, E_{a_1} , E_{b_3} and \widetilde{L}_{ab} is respectively 3, 1, 1 and 1; furthermore no other curve of Figure 1 intersects any curve of Ω_α . The situation for the curves of Ω_β is similar, after exchanging the roles of $E_{p(\alpha)}$ and $E_{p(\beta)}$.

Since $E_{b_3} \cong \mathbb{P}^1$, the points of E_{b_3} that do not lie on \widetilde{L}_{bc} or \widetilde{E}_{b_2} are parametrised by \mathbb{K}^* . Explicitly, the morphism $\pi' : X' \rightarrow \mathbb{P}^2$ is given locally by $(x, y) \mapsto (xy^4 : 1 : y)$, and in these coordinates, we define for any $\theta \in \mathbb{K}^*$ the point $q(\theta) \in E_{b_3} \subset X'$ that corresponds to $(\theta, 0)$. Any curve of Ω_α (respectively of Ω_β) passes through $q(\theta)$, for some $\theta \in \mathbb{K}^*$.

We assume that both Γ_1 and Γ_2 pass through the same point $q(\lambda) \in X'$, which is consistent with the fact that Γ_1 and Γ_2 have their maximum intersection at b . Blowing-up $q(\lambda)$, the exceptional curve $E_{q(\lambda)}$ intersects \widetilde{E}_{b_3} in one point, through which no curve of Ω_α or Ω_β passes. The remaining points of $E_{q(\lambda)}$ are parametrised by \mathbb{K} . Using the same coordinates as above, the blow-up of $q(\lambda) = (\lambda, 0)$ may be viewed as $(x, y) \mapsto (xy + \lambda, y)$, and the parametrisation associates to $\mu \in \mathbb{K}$ the point $r(\lambda, \mu)$, equal to $(\mu, 0)$.

Lemma 3.1. *For any pair $(\lambda, \mu) \in \mathbb{K}^* \times \mathbb{K}$, there exists a unique curve in Ω_α , that passes through $q(\lambda)$ and $r(\lambda, \mu)$. The same is true for Ω_β . The equations of the two curves are*

$$(2) \quad \begin{aligned} &\lambda^2 z(\alpha z - x)^3 + \alpha^2 xy^2(\mu(\alpha z - x) - \alpha \lambda y), \\ &\lambda^2 z(\beta z - x)^3 + \beta^2 xy^2(\mu(\beta z - x) - \beta \lambda y). \end{aligned}$$

Proof. This follows from a straightforward calculation, using the description of the blow-up in coordinates given above. \square

From now on, we fix $(\lambda, \mu) \in \mathbb{K}^* \times \mathbb{K}$, and denote by $\Gamma_1 \subset \Omega_\alpha$ and $\Gamma_2 \subset \Omega_\beta$ the two curves yielded by Lemma 3.1. Blowing-up the point $q(\lambda)$ on X' and then the point $r(\lambda, \mu)$, we obtain the birational morphism $\pi : X \rightarrow \mathbb{P}^2$ announced in the introduction. The situation on the blow-up of X' at $q(\lambda)$ and on the surface X is described in Figure 2.




Figure 2. The situation on the surface X . Two curves are connected by an edge if their intersection is positive (and here equal to 1). The positive intersections with Γ_1 and Γ_2 are in square brackets.

On the surface X , let R be the reducible curve which is the union of the 9 curves of self-intersection ≤ -2 of Figure 2 (the curves in grey).

Proposition 3.2. Fix some $i \in \{1, 2\}$. There exists a birational morphism $\eta_i : X \rightarrow \mathbb{P}^2$ that collapses the curves $R \cup \widetilde{\Gamma}_i$; it starts by collapsing $\widetilde{\Gamma}_i$ and then collapses the images of respectively \widetilde{L}_{ab} , \widetilde{E}_b , \widetilde{E}_{b_1} , \widetilde{E}_{b_2} , \widetilde{E}_{b_3} , $\widetilde{E}_{q(\lambda)}$, \widetilde{L}_{bc} , \widetilde{L}_{ac} , \widetilde{E}_a .

Then, $\eta_i(\widetilde{\Gamma}_{3-i})$ is a curve of \mathbb{P}^2 of degree 39, irreducible over the algebraic closure of \mathbb{K} , which has exactly one singular point. The morphism η_i is a minimal resolution of this curve, and is a (-1) -tower resolution of it (see Definition 2.1).

Proof. The curve $\widetilde{\Gamma}_i$ is a (-1) -curve (a smooth rational curve of self-intersection -1 , irreducible over the algebraic closure of \mathbb{K}). We may therefore collapse it and obtain a birational morphism $X \rightarrow Y$ where Y is smooth and projective. On Y , the image of \widetilde{L}_{ab} is a (-1) -curve so we may collapse it. Continuing with the images of $\widetilde{E}_b, \widetilde{E}_{b_1}, \dots, \widetilde{E}_a$ we obtain a birational morphism $\eta_i : X \rightarrow Z$ for some smooth rational projective surface Z (see Figure 3).




Figure 3. The decomposition of the birational morphism $X \rightarrow Z$ and the image of $R \cup \widetilde{\Gamma}_i$ on each surface.

Since X was obtained by blowing-up 10 points from \mathbb{P}^2 and η_i collapses 10 irreducible curves, we have $(K_Z)^2 = (K_{\mathbb{P}^2})^2 = 9$, so $Z \cong \mathbb{P}^2$.

Write $j = 3 - i$. Since $\widetilde{\Gamma}_j$ is not collapsed by η_i , the image $\eta_i(\widetilde{\Gamma}_j)$ is a curve. Its irreducibility follows from that of $\widetilde{\Gamma}_j$. Its degree can be calculated by computing its self-intersection after each of the 10 blow-downs. Since $R \cup \widetilde{\Gamma}_i$ is connected, its image by η_i is a single point. The curve $\widetilde{\Gamma}_j$ is smooth and intersects $\widetilde{\Gamma}_i$ in more than one point, hence η_i is a minimal resolution of $\eta_i(\widetilde{\Gamma}_j)$ and this curve has a unique singular point. Furthermore η_i is a tower resolution, as it collapses only one curve of self-intersection -1 . \square


4. Comparison of the curves $\eta_1(\widetilde{\Gamma}_2)$ and $\eta_2(\widetilde{\Gamma}_1)$

Proposition 3.2 shows that for any choice of $\alpha, \beta, \lambda \in \mathbb{K}^*$, $\alpha \neq \beta$, $\mu \in \mathbb{K}$, the complements of the two curves $\eta_1(\widetilde{\Gamma}_2)$ and $\eta_2(\widetilde{\Gamma}_1)$ are isomorphic. In this section, we distinguish the differences between the two curves.

Proposition 4.1. *The following are equivalent:*

- (1) *There exists an automorphism ψ of \mathbb{P}^2 that sends $\eta_1(\widetilde{\Gamma}_2)$ on $\eta_2(\widetilde{\Gamma}_1)$.*
- (2) *There exists an automorphism ψ' of X that leaves invariant every irreducible component of R and exchanges $\widetilde{\Gamma}_1$ and $\widetilde{\Gamma}_2$.*
- (3) *There exists an automorphism ψ'' of \mathbb{P}^2 that fixes a, b and c and permutes Γ_1 and Γ_2 .*
- (4) *$\mu = 0$ and $\alpha + \beta = 0$.*

Proof. Let us keep Diagram 1 in mind. The fact that η_1 (respectively η_2) is a minimal resolution of $\eta_1(\widetilde{\Gamma}_2)$ (respectively of $\eta_2(\widetilde{\Gamma}_1)$) and the assumptions made on the automorphisms above imply that ψ' may be constructed starting from ψ , as $\psi' = \eta_1^{-1}\psi\eta_2$. Similarly, the existence of ψ' implies that of ψ and ψ'' , constructed as $\psi = \eta_1\psi'\eta_2^{-1}$ and $\psi'' = \pi\psi'\pi^{-1}$. Finally, if ψ'' exists, then $\psi' = \pi^{-1}\psi''\pi$ exists.



It remains to prove that assertions (3) and (4) are equivalent. If ψ'' exists, then it is of the form $(x : y : z) \mapsto (x : \xi y : \theta z)$, for some $\xi, \theta \in \mathbb{K}^*$. Since ψ'' exchanges the curves Γ_1 and Γ_2 , it exchanges the points $p(\alpha) = (\alpha : 0 : 1)$ and $p(\beta) = (\beta : 0 : 1)$, which implies that $\alpha + \beta = 0$ and $\theta = -1$. Using the explicit equations of Γ_1 and Γ_2 , we find directly that $\mu = 0$. Conversely, if $\mu = 0$ and $\alpha + \beta = 0$ the automorphism $(x : y : z) \mapsto (x : y : -z)$ exchanges Γ_1 and Γ_2 . \square

Propositions 3.2 and 4.1 yield counterexamples to Conjecture 1.1, for any field \mathbb{K} that has more than two elements. Now, we study more intrinsically the curves $\eta_1(\widetilde{\Gamma}_2)$, $\eta_2(\widetilde{\Gamma}_1)$, without taking into account the plane embedding.

Proposition 4.2. *Neither $\eta_1(\widetilde{\Gamma}_2)$ nor $\eta_2(\widetilde{\Gamma}_1)$ is a curve of type I.*

If $\mu = 0$, the curves $\eta_1(\widetilde{\Gamma}_2)$ and $\eta_2(\widetilde{\Gamma}_1)$ are isomorphic.

For any field \mathbb{K} with more than two elements there exist values of α, β, μ for which the curves $\eta_1(\widetilde{\Gamma}_2)$ and $\eta_2(\widetilde{\Gamma}_1)$ are not isomorphic.

Proof. Denote by p_1 (respectively p_2) the morphism $\widetilde{\Gamma}_1 \rightarrow \eta_2(\widetilde{\Gamma}_1)$ (respectively $\widetilde{\Gamma}_2 \rightarrow \eta_1(\widetilde{\Gamma}_2)$) obtained by restriction of η_2 (respectively η_1). The singular curves $\eta_1(\widetilde{\Gamma}_2)$ and $\eta_2(\widetilde{\Gamma}_1)$ are isomorphic if and only if there is an isomorphism $\rho : \widetilde{\Gamma}_1 \rightarrow \widetilde{\Gamma}_2$ that is compatible with p_1 and p_2 . Furthermore the singular curves are of type I if and only if the morphisms p_i are injective.

The ramified form of the morphism p_1 consists of the point $\widetilde{L}_{ab} \cap \widetilde{\Gamma}_1$ and the form of degree 8 on $\widetilde{\Gamma}_1 \cong \mathbb{P}^1$ obtained by intersecting $\widetilde{\Gamma}_1$ with $\widetilde{\Gamma}_2$. Taking some coordinates

$(u : v)$ on $\widetilde{\Gamma}_1 \cong \mathbb{P}^1$, the morphism $\widetilde{\Gamma}_1 \rightarrow \Gamma_1 \subset \mathbb{P}^2$ obtained by restriction of π is the following:

$$(u : v) \mapsto (v^4 \lambda^5 \alpha : (u + \mu v)(\alpha(u + \mu v)^2 u - \lambda^4 v^3) : v(u + \mu v)^2 \lambda u \alpha).$$

The point $(1 : 0)$ is sent on b , the point $(\mu : -1)$ is sent on a and the point $(0 : 1)$ corresponds to $\widetilde{\Gamma}_1 \cap \widetilde{L}_{ab}$. Replacing the parametrisation in the equation of Γ_1 we find 0, and replacing it in the equation of Γ_2 , we find

$$-\lambda^{10} \alpha(\alpha - \beta)(u + \mu v)^2 v^7 \left(\sum_{i=0}^7 c_i \cdot u^i v^{7-i} \right),$$

where c_0, \dots, c_7 are as follows:

$$\begin{aligned} c_0 &= 3\alpha^2\beta^2, & c_4 &= -\alpha\beta\mu(8\lambda^4\beta - 7\alpha\beta\mu^3 + 6\lambda^4\alpha), \\ c_1 &= 13\alpha^2\beta^2\mu, & c_5 &= -\alpha\beta\mu^2(3\lambda^4\alpha - \alpha\beta\mu^3 + 7\lambda^4\beta), \\ c_2 &= 22\alpha^2\beta^2\mu^2, & c_6 &= \lambda^4(\lambda^4(\alpha\beta + \alpha^2 + \beta^2) - 2\alpha\beta^2\mu^3), \\ c_3 &= -3\alpha\beta(\lambda^4(\alpha + \beta) - 6\alpha\beta\mu^3), & c_7 &= \lambda^8\beta^2\mu. \end{aligned}$$

The intersection number of Γ_1 and Γ_2 is 16; the intersections at a and a_1 correspond to the factor $(u + \mu v)^2$ and the intersections at $b, b_1, b_2, b_3, q(\lambda), r(\lambda, \mu)$ correspond to v^6 . Thus, the form of degree 8 on $\widetilde{\Gamma}_1$ corresponding to the intersection of $\widetilde{\Gamma}_1$ and $\widetilde{\Gamma}_2$ is $F_1 = v \sum_{i=0}^7 c_i \cdot u^i v^{7-i}$. Since F_1 vanishes at the point $(1 : 0)$ and $(0 : 1)$ corresponds to $\widetilde{\Gamma}_1 \cap \widetilde{L}_{ab}$, the map p_1 is not injective and $\eta_2(\widetilde{\Gamma}_1)$ is not of type I.

For $p_2 : \widetilde{\Gamma}_2 \rightarrow \eta_1(\widetilde{\Gamma}_2)$, the situation is similar. We find a form F_2 , that is equal to F_1 , after exchanging α and β . As above, we see that $\eta_1(\widetilde{\Gamma}_2)$ is not of type I. Finally, the two singular curves are isomorphic if and only if there exists an isomorphism of \mathbb{P}^1 that fixes $(0 : 1)$ and sends F_1 on F_2 (we say in this case that F_1 and F_2 are equivalent). If $\mu = 0$, the identity suits, since each c_i becomes symmetric with respect to α and β . If $\mu \neq 0$, this is not the case. If $\text{char}(\mathbb{K}) \neq 2$, choosing $\alpha = 1, \beta = 2, \lambda = \mu = 1$, we can compute that F_1 and F_2 are not equivalent. If $\text{char}(\mathbb{K}) = 2$, there is considerable simplification of the terms, and we find that if $\lambda = \mu = 1, \alpha \neq \beta$, then F_1 and F_2 are not equivalent. \square

The proof of Theorems 1 and 2 now follows directly from Propositions 3.2, 4.1 and 4.2.

References

- [1] *S. Abhyankar, T. Moh*, Embeddings of the line in the plane, *J. reine angew. Math* **276** (1975), 148–166.
- [2] *M. H. Gizatullin, V. I. Danilov*, Automorphisms of affine surfaces. I, *Izv. Akad. Nauk SSSR Ser. Mat.* **39** (1975), no. 3, 523–565, 703.
- [3] *S. Iitaka*, On logarithmic Kodaira dimension of algebraic varieties, *Complex analysis and algebraic geometry*, Iwanami Shoten, Tokyo (1977), 175–189.

- [4] *T. Kishimoto*, Projective plane curves whose complements have logarithmic Kodaira dimension one, Japan. J. Math. (N.S.) **27** (2001), no. 2, 275–310.
- [5] *H. Kojima*, On the logarithmic plurigenera of complements of plane curves, Math. Ann. **332** (2005), no. 1, 1–15.
- [6] *M. Miyanishi*, Noncomplete algebraic surfaces, Lect. Notes Math. **857**, Springer-Verlag, Berlin-New York 1981.
- [7] *M. Miyanishi*, Open algebraic surfaces, CRM Monogr. Ser. **12**, American Mathematical Society, Providence, RI, 2001.
- [8] *I. Wakabayashi*, On the logarithmic Kodaira dimension of the complement of a curve in \mathbb{P}^2 , Proc. Japan Acad. Ser. A Math. Sci. **54** (1978), 157–162.
- [9] *H. Yoshihara*, On plane rational curves, Proc. Japan Acad. Ser. A Math. Sci. **55** (1979), no. 4, 152–155.
- [10] *H. Yoshihara*, Rational curve with one cusp, Proc. A.M.S. **89** (1983), 24–26.
- [11] *H. Yoshihara*, On open algebraic surfaces $\mathbb{P}^2 - C$, Math. Ann. **268** (1984), 43–57.
- [12] *H. Yoshihara*, Projective plane curves and the automorphism groups of their complements, J. Mat. Soc. **37** (1985), no. 1, 87–113.
- [13] *H. Yoshihara*, Rational curve with one cusp, II, Proc. A.M.S. **100** (1987), 405–406.

Université de Grenoble I, UFR de Mathématiques, UMR 5582 du CNRS, Institut Fourier, BP 74,
38402 Saint-Martin d'Hères, France
e-mail: Jeremy.Blanc@fourier.ujf-grenoble.fr

Eingegangen 19. Februar 2008